

# METHANDROSTENOLONE 10 mg MUSCLE Tablets Via Oral

## FORMULA:

### Each tablet contains:

Methandrostenolone.....1

00mg.

Excipients.....

c.s.p.

## MECHANISM OF ACTION

Methandrostenolone is characterized by having anabolic and androgenic properties. Promotes protein synthesis because it supports muscle growth. This effect is sustained by a positive nitrogen balance and an improved feeling of well-being. The calcium balance is also positively influenced as it favors the deposition of calcium in the bones. The biological half-life is approximately 3 to 4 hours. It is eliminated in the urine.

## INDICATIONS

Methandrostenolone is indicated in the treatment of all diseases and conditions in which an anabolic effect (increased protein synthesis) in the muscles and a general strengthening effect may be needed.

## POSOLOGY

Recommended dose: 20mg. to 60mg. daily in 3 or 4 doses. Beginners only need 15mg to 20mg. daily. Doses taken by professionals range from 15 mg. to 40mg. per day, rarely needing to reach 50 mg per day. Using more than 50 mg per day does not make sense since beyond these levels the increase in effect is not proportional to the increase in dose but much less. Its use in women is not recommended because it is highly androgenic and can cause virilization.

## CONTRAINDICATIONS

Hypersensitivity to methandrostenolone. Severe liver failure. In pre-pubertal individuals or in an aggressive state. Pregnancy, lactation.

## ADVERSE EFFECTS

Las siguientes reacciones requieren acompañamiento medico: Virilización e irregularidades menstruales en mujeres, irritabilidad de laveziga, ginecomas

## ADVERSE REACTIONS

There may be menstrual irregularities in women, bladder irritability, gynecomastia, Anaphylaxis, edema, erythrocytosis, gastrointestinal infection, hypercalcemia, polycythemia, androgenital alopecia, seborrhea and acne in men and women. Prostatic hypertrophy and increased sexual desire, cholestatic jaundice, excitability, hypercholesterolemia, hypercalcemia.

edema. In women virilization. In men, priapism, oligosperm, hirsutism, reduced libido, decreased testicular mass, functional impotence. Other adverse reactions are constipation, nausea, diarrhea, infection, libido changes, stomach pain, sleep difficulties, headaches, anxiety, depression, generalized paresthesia, skin rash.

## PRECAUTIONS AND WARNINGS

Methandrostenolone is associated with hepatotoxicity and liver function should be monitored during treatment. It should not be administered in patients with hepatic insufficiency.

## INTERACTIONS

Like testosterone and other androgens and anabolic steroids, they potentiate the activity of some drugs and consequently their toxicity. Among them, cyclosporine, antidiabetics, levothyroxine, anticoagulants such as warfarin stand out. Resistance to neuromuscular blocking agents has also been described.

## USE RESTRICTIONS

Professional use.

## OVERDOSE

There is no specific antidote, therefore supportive measures should be taken, such as maintaining a patent airway and monitoring cardiovascular function. In all cases, resort to the Medical Emergency Center

## PRESENTATION

Box containing bottle x 100 tablets

Store at room temperature between 15°C - 30°C.  
Keep out of the reach of children.