

**PARABOLON BLEND**  
**TREMBOLONE SALTS mg/ml. MUSCLE**  
**INJECTABLE OIL SOLUTION / INTRAMUSCULAR WAY**

**Composition of this medicine**

Every 1 ml. of injectable Oily Solution contains:

Trembolone acetato.....	50 mg.
.Trembolone hexahydroxybenzi carbonato.....	50 mg.
Trembolone enantato.....	100 mg.
Excipientes.....	c.s.p.

**What are Trenbolone salts?** It is a very potent steroid mix drug that contains three of the most active rebolone actives including Trenbolone Acetate, Trenbolone Enanthate and Trenbolone Hexahydrobenzylcarbonate which belong to the Androgen group. The presence of two long-base versions of enanthate ester and hexahydrobenzylcarbonate makes the steroid relatively slow acting. Trenbolone is basically in charge of developing and maintaining the body in recovery from debilitating catabolic processes and helping various systems, including the muscle.

**How does this medicine work?**It works by activating its highly selective anti-catabolic properties, helping to maintain protein levels in the body, increasing erythropoiesis and raising the state of conservation and development of the muscular system, as well as regulating fats.

**What is this medicine for?**For the treatment of all conditions related to general commitment disorders, it is an ideal drug for states of convalescence, for burnt or weakened patients and, above all, to prevent muscular atrophy. It also has hormonal and functional activity in man. It is also used to treat other debilitating and debilitating chronic diseases.

**Who should be careful when using this medication?**Those with heart disease, high blood pressure, tumors, breast and prostate cancer, prostatic hyperplasia, behavioral disorders, liver problems, poor kidney function, imbalanced calcium in the blood, thyroid, acne, excessively oily skin, respiratory conditions severe or chronic. Especially in women, due to its hormonal action, it is prohibited and contraindicated. Also children, adolescents and seniors. Pregnant and lactating women as it is potentially risky. It should not be administered to those who receive other immunodebitating treatments, they must necessarily consult their doctor.

**THERAPEUTIC ACTION:** Anticatabolic, malnutrition, cachexia, physical decay, weakness and general compromise, burnt recovery, prevention of muscle atrophy, selective fat regulator, metabolic cycle activator, increases lean mass, increases hardness, definition and muscle strength, increases erythropoiesis

**POSOLOGY**

Recommended dose of trenbolonA salts 150mg or according to medical criteria.

**ACTION MECHANISMS**

Trenbolone is a substance that belongs to the group of non-stilbene xenobiotics, therefore it is a synthetic hormone derived from Nortestosterone, thus being a modified derivative of Nandrolone and differs from it by including two double-characteristic bonds in the Carbons. 9 and 11, which inhibits its aromatization so characteristic of this type of hormones (9-en), increasing its androgenic affinity and slowing down its metabolism. At the C9 and C10 bond level, it inhibits Estrogenic activity, although it seems that this substance itself is not metabolically canceled since these compounds are necessary for it to become Estrogen, so the final steroid (Trenbolone) is 3 times more powerful in its action anabolic as androgenic than nandrolone base. Since Trenbolone is not aromatized in the body, it is no longer Estrogenic, presenting greater affinity for the Progesterone receptors, therefore the benefits are demonstrated in that the side effects due to the aromatization of Estrogen are practically not present and because the already modified Trenbolone continues to act as a highly potent muscle mass, strength and definition drug much more so than Testosterone. But the most striking thing is that due to its steroidal characteristics, Trenbolone reaches a greater amount of time within the body after its administration, in addition to the strong impact it has on endogenous testosterone, it is safer to use drugs such as HCG, after finishing a treatment cycle since without the use of this type of drugs the hormonal leveling could last a long time

At the testicular level of endogenous gonadotropins.

**ADVERSE BIOLOGICAL EFFECTS OF TREMBOLONE IN GENERAL:**Woman: Severe virilization, Larynx: Growth and elongation of the vocal cords, Skin: Stimulates the production of fat and Acne, Prostate: stimulates its growth (HBP), Brain: temporary inhibition of respiratory centers especially at the time of administration (Tren Cough ). Sexual and reproductive disorders due to high levels of Prolactin, Increased aggressiveness (rare)

**WARNINGS AND CAUTIONS:**It should not be used for long periods. Do not administer to patients with other treatments. Do not consume alcohol simultaneously. Do not exceed doses. Keep out of the reach of children. Administer with caution to patients who are allergic or sensitive to this medication or its components. **WARNING!** Warnings for athletes have NOT been described, but it is reported that this medicine contains a component that in a drug test can give the result as POSITIVE (+).

**ADVERSE REACTIONS:** The appearance of adverse reactions can be: Virilization in women (increased tone of voice, hirsutism, clitoral hypertrophy, menstrual irregularities, amenorrhea, thickening of the skin and others) In adults Acne. Moderate polycythemia and obstruction due to Prostatic Hyperplasia, Liver Diseases (Cholestatic Jaundice), Hepatic Carcinoma, Prostate Cancer Mild Hydrosaline Retention, Arterial Hypertension. In small men and slight feminizing effects, (mild gynecomastia, decreased libido, probable erection disorders and impotence of the Adrenal alteration type: | end the administration cycle. Hormonal alterations in the Thyroid, Nephropathy and atuteration of the adrenal glands .

**DRUG INTERACTIONS:**Barbiturates: Microsomal enzyme inducers, alter clearance, Amiodarone, Antibiotics, Phenytoin-Oral Anticoagulants: Coumarinics with increased activity (check dose), INR and Prothrombin control, edema formation (mild). Do not consume alcohol simultaneously.

**USE RESTRICTIONS:**Especially in women due to its hormonal action it is prohibited and contraindicated. Also children, adolescents and the elderly. Pregnant and lactating women as it is potentially risky.

**OVERDOSE**

-Turn quickly in Paraguay to the Toxicology Center of the Hospital for Medical Emergencies .

**PRESENTATION**

Box containing 1 vial x 10 ml (multidose)  
Box containing 5 and 10 vials of 1 mL

**Keep at room temperature (15°C- 30°C)**  
**Keep out of the reach of children.**