

**MASTERON-MUSCLE  
DROSTANOLONE  
PROPIONATE 100 mg/mL**  
Injectable oily solution  
Deep intramuscular routea

**FORMULA**

Each mL of injectable oily solution contains:

Drostanolona

Propionato.....100m

g.

Excipients.....

..... C.S.

**MECHANISM OF ACTION**

The mechanism of action of Drostanolone Propionate is based on the following principles:- It acts by favoring the increase in strength and muscle mass through various physiological mechanisms by increasing nitrogen retention.- It is myotropic because it acts in the cytoplasm of the muscle cell, promotes the release of the alpha reductase enzyme in the nucleus, allowing the RNA to take advantage of the amino acids and proteins (nitrogen) from the diet to transform them into muscle tissue.- Likewise, it has the action of retaining calcium, phosphorus, potassium and chlorides. This action contributes to a greater development of the bones, also constituting a growth factor, as long as the recommended dosages are maintained. These construction processes of mineralization and consolidation also allow bone regeneration in cases of rickets and osteomalacia.- Appetite stimulation through metabolic regulation.- It favors the absorption of the nutrients acquired by the diet or supplied as a supplement.

**DOSAGE AND INSTRUCTIONS FOR USE**

The recommended route is deep intramuscular, according to the following doses:1 mL for every 90 kg of weight. Although the duration of treatment depends on the discretion of the doctor,The following dosing models are recommended:- Do not administer intravenously.- Do not mix in the same syringe or container with any other substance foreign to the product.- Keep the indications of asepsis and antisepsis before and during the application of the product.- To obtain an adequate effect, it is important to maintain an adequate intake of proteins, calories, vitamins, iron and copper.- A slight swelling may appear at the inoculation site, which disappears after a few days without the need for treatment- Keep out of the reach of children and pets.- Do not smoke or eat while handling the product.

**INDICATIONS**

Drostanolone Propionate has been used in most cases by athletes in phases of pre-competition, cutting or simply to avoid water retention and estrogen spikes. For mass and volume Drostanolone Propionate is more effective in cycles with other drugs such as the anabolic Stanozolol, Testosterone Propionate and others. Specific indications include:- Greater development of bone structure and increase in muscle mass.- Better and greater protein synthesis, mineral retention, improvement of appetite and feed conversion.- As an adjuvant in wasting or debilitating diseases that cause weight loss, states

**MECHANISM OF ACTION**

The mechanism of action of Drostanolone Propionate is based on the following principles:- It acts by favoring the increase in strength and muscle mass through various physiological mechanisms by increasing nitrogen retention.- It is myotropic because it acts in the cytoplasm of the muscle cell, promotes the release of the alpha reductase enzyme in the nucleus, allowing the RNA to take advantage of the amino acids and proteins (nitrogen) from the diet to transform them into muscle tissue.- Likewise, it has the action of retaining calcium, phosphorus, potassium and chlorides. This action contributes to a greater development of the bones, also constituting a growth factor, as long as the recommended dosages are maintained. These construction processes of mineralization and consolidation also allow bone regeneration in cases of rickets and osteomalacia.- Appetite stimulation through metabolic regulation.- It favors the absorption of the nutrients acquired by the diet or supplied as a supplement.

**DOSAGE AND INSTRUCTIONS FOR USE**

The recommended route is deep intramuscular, according to the following doses:1 mL for every 90 kg of weight. Although the duration of treatment depends on the discretion of the doctor,The following dosing models are recommended:

- Do not mix in the same syringe or container with any other substance foreign to the product.
- Keep the indications of asepsis and antisepsis before and during the application of the product.
- To obtain an adequate effect, it is important to maintain an adequate intake of protein, calories, vitamins, iron and copper
- A slight swelling may appear at the inoculation site, which disappears after a few days without the need for treatment.
- Keep out of the reach of children and pets.
- Do not smoke or eat while handling the product