

MUSCLELABS CUT STACK

TRENBOLONE ENANTHATE 50 mg/ml
TESTOSTERONE CYPIONATE 100 mg/ml
DROSTANOLONE PROPIONATE 50 mg/ml

FORMULA:

Each 1 ml. contains:

Drostanolone Propionate.....	100mg.
Testosterone propionate.....	50mg.
Trenbolone Acetate.....	100mg.
Excipient.....	c.s.p.

PROPERTIES:

Cutstack is a mixture of Testosterone Cypionate + Dronastonolone Propionate + Trenbolone Enanthate. It is generally used within cycles and is very effective. Thanks to Cutstack, the body can retain water for longer. Shows the result in a short time and with short VM. For this reason, it is an effective combination for athletes who are concerned about anti-doping testicles.

INDICAÇÕES:

Cut Stack is generally used within cycles, and is very effective. Thanks to Cutstack, the body can maintain water for longer. She showed the result in a short period of time and with short MV. For this reason, it is an effective combination for athletes.

CONTRAINDICATIONS

Contraindicated if you have hypersensitivity to these substances and you should never use any anabolic steroid..

DOSE:

- Twice a week, for a period of 4 weeks.

PRECAUTIONS AND WARNINGS:

Do not treat palliative breast cancer, do not continue therapy if the condition persists after 3 months or hypocalcemia may occur at any stage of the disease. Androgen therapy for women, in addition to being short-term, can produce virilization, especially vocal and hairy. Androgen therapy is not advice to improve sports rest. In children, this medication should be used with caution with adverse effects on the bone maturation process, which may accelerate the production of compensatory gains without linear growth. When patients over 50 years of age receive androgen treatment, their risk of prostate enlargement or the development of prostate cancer increases. For this reason, it is necessary to perform prostate and blood tests before prescribing androgens to homens de este idade. During prolonged treatment, oligospermia, azoospermia or reduced sperm function may occur. Resulting in possible infertility, with spontaneous remission after discontinuation of treatment. Patients with benign prostatic hypertrophy may develop acute urethral obstruction, requiring immediate discontinuation of the medication. Urinary calcium levels should be determined frequently in women such as metastatic breast carcinoma treated with testosterone. During the treatment of pre-pubertal minors, bone x-rays must be performed every 6 months. In patients with acute intermittent porphyria, androgens may precipitate attacks of that condition. It is generally preferable to start treatment with full doses and subsequently adjust them to individual doses. characteristics.

OVERDOSE

Frequent doses may cause symptoms of slow remission, as it is a long-acting medication. It is recommended to immediately discontinue the medication.

RECOMMENDATIONS:

-Using a wet needle or syringe may cause problems with the solution, so it does not affect the effectiveness of the medication. If crystals or bubbles form, gently heating, rubbing between your hands or shaking can dissolve the crystals.

BOX CONTAINS:

- Bottle with 10mL content

Store at a temperature below 25°C in a dry place.
and out of reach of the children.